



IFSO Melbourne 2024

Edo Aarts





IMPACT STUDY

**Preoperative Micronutrient
Supplementation**



Edo Aarts, MD, PhD



Dr. Johannes Sander





ETHICON

Medtronic

fitforme

dockx

Allurion
KLINIEK




SCHÖN KLINIK
Hamburg Eilbek

WEIGHTWORKS
specialist in maagverkleiningen



General



WLS Start enthält pro Tagesdosis			
Vitamine	Verbindung	Menge	RI**
A	Retinylpalmitat	1000 mcg RE/ER	125%
-	Beta-Carotin	0,2 mg	-
B1*	Thiaminhydrochlorid	5,5 mg	500%
B2*	Riboflavin	0,7 mg	50%
B3*	Nicotinamid	8 mg NE	50%
B5*	Calcium-D-pantothenat	6 mg	100%
B6*	Pyridoxal-5'-phosphat	0,7 mg	50%
B8*	D-Biotin	50 mcg	100%
B9/B11	Pteroylmonoglutaminsäure	600 mcg	300%
B12*	Methylcobalamin	100 mcg	4000%
C*	L-Ascorbinsäure	250 mg	313%
D	Cholecalciferol	75 mcg	1500%
E	D-a-Tocopherolsäuresuccinat	12 mg a-ET/TE	100%
Mineralstoffe			
Eisen*	Eisenfumarat	24 mg	172%
Zink	Zinkcitrat	22,5 mg	225%
Kupfer*	Kupfergluconat	2,25 mg	225%
Chrom	Chrom(III)-Chlorid	80 mcg	200%
Jod*	Kaliumiodid	150 mcg	100%
Selen	Natriumselenat	100 mcg	182%
Mangan*	Mangancitrat	2 mg	100%
Molybdän	Natriummolybdat	50 mcg	100%

Inhaltsstoffe:
Vitamine, Mineralien, Magnesiumcitrat, Kapsel (Hypromellose), Füllstoff (mikrokristalline Cellulose), Stabilisator (Siliciumdioxid), Trennmittel (Magnesiumstearat).

** RI (reference intake) = Referenzmenge nach der Verordnung (EU) 1169/2011.


Chargennummer: siehe Packungsseite und/oder Blister
Mindestens haltbar bis Ende: siehe Packungsseite und/oder Blister
Nettogewicht: 41 gram



Setup



IMProving blood levels before bAriatriC surgery in Three months: Effects of a specialized multivitamine



120 patients included

3 months

Bloodcheck 1


Bloodcheck 2

....

Surgery



Results




Decrease in ferritin and B12 **not clinically relevant:**

- Ferritin: decrease blood loss during menstruation
- B12: decrease only in people with high initial value



Results





Conclusion



WLS Start is effective in improving blood serum values

WLS Start is effective in decreasing deficiencies

3 Months of use before surgery appears safe