

# Stomach Viability after Hiatal Hernia Reduction Containing the Gastric Sleeve


**Dr. Thelma E. Encarnación Santana, Dr. Isabel Molina Céspedes, Dr. Alberto Michel, Dr. Pamela Huerta, Dr. Nadia Nazrala**



I have no potential conflict of interest to report




- 56 – year- old woman with **gastric sleeve surgery 8 years prior.**



- Symptoms of **GERD** in the last 6 months and **weight regain.**
- Upper GI endoscopy revealed a **large hiatal hernia** and pouch migration.
- The patient was scheduled to surgery to perform Hiatal hernia repair and RYGB


## Stomach Viability after Hiatal Hernia Reduction Containing the Gastric Sleeve



# Postoperative Follow Up

- Discharged without complications after 2 days.
- Asymptomatic on subsequent visits
- After **3 months** we performed the RYGB

Stomach Viability after Hiatal Hernia Reduction Containing the Gastric Sleeve



# Conclusion

- **21-34%** of LSG will have revisional surgery because Hiatal Hernia or GERD.
- Tools like endoscopy, Indocyanine green, etc are a **MUST** in revisional surgery.
- It is prudent to wait and do surgery in a second time if risk to the patient are observed during revisional surgery.



Felsenreich D, et al. OBES SURG (2017) 27:3092–3101.  
Zundel N, et al. Laparoscopic Sleeve Gastrectomy: Technique and Outcomes. The  
ASMBS Textbook of Bariatric Surgery. 2020.

Thanks for your attention

