


GLP-1RAs & MBS:

How to Rescue, Where Hope Returns

NO Semaglutide or Tirzepatide Before Surgery




Cleveland Clinic Date
N=4661

T2DM, RYGB (29%)

No T2DM, RYGB (33%)

*Under Peer Review.
Please do not share.*

NO Semaglutide or Tirzepatide Before Surgery



Cleveland Clinic Date
N=4661

T2DM, Sleeve (20%)


No T2DM, Sleeve (24%)

T2DM, RYGB (29%)

No T2DM, RYGB (33%)


*Under Peer Review.
Please do not share.*

Preop Weight Change After Medication Initiation



*Under Peer Review.
Please do not share.*

Stopped Semaglutide or Tirzepatide Before Surgery for Side Effects




Cleveland Clinic Data
N=61

20-30%
weight loss

*Under Peer Review.
Please do not share.*

Ineffective Semaglutide or Tirzepatide Before Surgery




Cleveland Clinic Data
N=207

T2DM, RYGB (27%)

No T2DM, RYGB (36%)

*Under Peer Review.
Please do not share.*

Ineffective Semaglutide or Tirzepatide Before Surgery



Cleveland Clinic Date
N=207

T2DM, Sleeve


No T2DM, Sleeve

T2DM, RYGB

No T2DM, RYGB

*Under Peer Review.
Please do not share.*

Ineffective Semaglutide or Tirzepatide Before Surgery



Cleveland Clinic Date
N=207

T2DM, Sleeve X

No T2DM, Sleeve

T2DM, RYGB

No T2DM, RYGB

*Under Peer Review.
Please do not share.*